Focusing of a Parallel Beam to Form a Point in the Particle Deflection Plane

Ivar Ekeland[1], Roger Temam[2], Jeffrey Dean[2], David Grove[1], Craig Chambers[2], Kim B. Bruce[2], and Elsa Bertino[1]

[1] Princeton University, Princeton NJ 08544, USA
[2] Université de Paris-Sud, Laboratoire d'Analyse Numérique, Bâtiment 425, F-91405 Orsay Cedex, France

Abstract. The abstract should summarize the contents of the paper in at least 70 and at most 150 words; neither too long nor too short but to the point!

1 Fixed-Period Problems: The Sublinear Case

Of course this text has no scientific relevance but should be used as a practical example of the Springer layout specifications for *mathematics books* [1]. We begin now the search for periodic solutions to Hamiltonian systems. All this will be done in the convex case; that is, we shall study the boundary-value problem

$$\begin{aligned} \dot{x} &= JH'(t,x) \\ x(0) &= x(T) \end{aligned}$$

with $H(t,\cdot)$ a convex function of x, going to $+\infty$ when $\|x\| \to \infty$.

2 Equations

The following equation is the compiled result of using the command `\vec` and using the newly defined command `\umu`:

$$\boldsymbol{A} = \mu y = 50\,\mu\mathrm{m}\,. \tag{1}$$

2.1 Sub-equations

As you have seen above, equations are numbered automatically when the equations are defined with `\begin{equation}` (but not when `$$` is used). Here you will find an example for the automatic sub-numbering of equation arrays, using the style `subeqnar.sty`:

$$a = c + d\,, \tag{2a}$$

$$e = f - d\,. \tag{2b}$$

In order to refer to the equations within the main text you must simply label your equations (within the equation's environment) and quote the labels within the text environment. Upon running TEX you should receive the corresponding subnumbers, e.g. (2a) and (2b).

No Sub-numbering. If you want to suppress the sub-numbering of an equation array you use the original `\begin{eqnarray}` environment, put `\nonumber` at the end of the line that is to have no number, and set the sub-counter to zero:

$$\begin{aligned} a &= c + d\,, \\ e &= f - d\,. \end{aligned} \tag{3}$$

3 Figures

If your figures are available as electronic files it is advisable to convert them to eps format and include them directly into the text with the help of the graphicx package (see the following example in Fig. 1)



Fig. 1. Example of an electronically included eps figure

In order to mark the desired amount of space for a (centered) figure that is to be pasted into the manuscript manually we would ask you to provide a frame showing the size and position of the figure. This is done by using the commands `\centering\noindent\mpicplace{width cm}{height cm}`.

If there is enough space left ($> 3\,\mathrm{cm}$) you are welcome to set the figure caption at the right-hand side of the figure using the commands `\sidecaption\mpicplace{width cm}{height cm}`. For further instructions, e.g. on the structure and layout of the caption, see Fig. 2 und 3.



Fig. 2. General description. (**a**) The 'name' of the sub-figure should be set boldface and in round brackets. (**b**) In general the last sentence of a figure caption should not end with a full stop



Fig. 3. This could show a figure consisting of different types of lines to describe individual aspects. These descriptive lines (*dotted line*) should be set italic and put into round brackets as shown in this sample figure caption. In general the last sentence of a figure caption (*straight line*) should not end with a full stop

4 Tables

In *Springer mathematics books*, tables as well as equations and figures should be centered. Table captions should be treated in the same way as figure legends, except that the table captions appear *above* the tables. Overwide tables should be reduced to the page width, if possible, or exceed the type

area by a maximum of 5 mm. Please check the TEX coding of the following tables, (Table 1 and 2), and use them as an example for setting your own tables.

Table 1. Critical N values

$M_\odot$	β_0	T_{c6}	γ	N_{crit}^{L}	N_{crit}^{Te}
30	0.82	38.4	35.7	154	320
60	0.67	42.1	34.7	138	340
120	0.52	45.1	34.0	124	370

Table 2. Please write your table caption here. Multi-line captions will automatically be set flush left, while single-line captions will be centered

Nominal dimension (mm)	Angle	Tolerance (µm)	Evaluation factor
10^1	1°	Untoleranced dimension	$c_1 = 0$
10^{-1}	1′ [a]	101–200	$c_1 = 1$
10^{-2}	1″ [b]	40–100	$c_1 = 2$
10^{-3}		< 50	$c_1 = 3$

[a] One minute of arc.
[b] One second of arc.

5 Lists

We have redefined the *itemize* environment (labelitemi) so that you will receive bullets instead of dashes to introduce the individual items. We think that this way the list

- is clearer
- looks better
- is more noticeable

References

1. Holzwarth F., Lenz J. et al. (1998) 1readme. Further Details on Layout and LATEX code. Springer, Berlin Heidelberg

2. Turner, J. S. (1974) Double–diffusive phenomena. Ann. Rev. Fluid Mech. **6**, 37–56
3. Turner, J. S. (1968) Double–diffusive phenomena. Ann. Rev. Fluid Mech. **33**, 168–200
4. Normand, C., Pomeau, Y. (1977) Convective instability: a physicist's approach. Rev. Mod. Phys. **49**, 581–623

Index

abstract 1

boundary-value problem 1

Hamiltonian systems 1